



HisCap 6FF

Index

1. Product Description.....	1
2. Purification Procedure.....	2
3. Cleaning-in-Place.....	4
4. Regeneration.....	4
5. Troubleshooting.....	4
6. Related Products.....	5

1. Product Description

Ni NTA Beads 6FF is based on a highly cross-linked 6% agarose with the same ligand as Ni NTA Beads (See Fig 1.), see table 1. In addition to being able to tolerate harsh reagent conditions (see Table 2), **Ni NTA Beads 6FF** is more stable due to its pressure-resistant matrix, which can withstand pressures up to 0.3 MPa, making the product is more suitable for industrial large-scale protein purification, which can be achieved at relatively high flow rates for the purification of target proteins.

HisCap 6FF is one of a range of prepacked, ready-to-use columns for affinity chromatography. It is packed with 1ml and 5ml of **Ni NTA Beads 6FF**. Five different packing sizes are available. **HisCap 6FF** has the standard interface and can be adapted to all kinds of chromatography system, such as ÄKTA. It is fast, simple and easy operation.




Fig.1. The chemical structures of Ni NTA Beads 6FF

Table 1. Characteristics of Ni NTA Beads 6FF

Item	Description
Matrix Spherical	Highly cross-linked 6% agarose
Static Binding Capacity	>40 mg 6×His-tagged protein/ml medium
Particle size	45-165 μm
Maximum Pressure	0.3 MPa, 3 bar
Storage Solution	1×PBS containing 20% ethanol
Storage Temperature	4-30°C

Table 2. Chemical compatibilities for Ni NTA Beads 6FF

Reagent	Stability
Reductants	5 mM DTE
	1mM DTT
	20 mM β-mercaptoethanol
	5 mM TCEP
Denaturants	10 mM reduced glutathione
	8 M urea 6 M Gua-HCl
Detergent	2% Triton™ X-100 (nonionic)
	2% Tween™ 20 (nonionic)
	2% NP-40 (nonionic)
	2% cholate (anionic)
	1% CHAPS (zwitterionic)





Table 2. Chemical compatibilities for **Ni NTA Beads 6FF** (Continued table)

Reagent	Stability
Other additives	500 mM imidazole 20% ethanol 50% glycerol 100 mM Na ₂ SO ₄ 1.5 M NaCl 1 mM EDTA 60 mM citrate

2. Purification Procedure

2.1 Buffer Preparation

The basic principle of the following recommended buffer and other buffer is low concentration of imidazole in Lysis and wash buffer and high in elution buffer. Water and chemicals used for buffer preparation should be of high purity. It is recommended to filter the buffers by passing them through a 0.22 µm or 0.45 µm filter before use. **Ni NTA Beads 6FF** can be used for the his-tagged protein purification under native conditions and denaturing conditions, which need different buffer. See table 3, table 4 and table 5.

Table 3. Recommended buffer for his-tagged protein purification under native conditions

Name	Volume	Ingredient
Lysis Buffer	1 L	50 mM NaH ₂ PO ₄ (7.80 g NaH ₂ PO ₄ ·2H ₂ O) 300 mM NaCl (17.54 g NaCl) 10 mM imidazole (0.68 g imidazole) Adjust the buffer pH to 8.0 with NaOH solution.
Wash Buffer	1 L	50 mM NaH ₂ PO ₄ (7.80 g NaH ₂ PO ₄ ·2H ₂ O) 300 mM NaCl (17.54 g NaCl) 20 mM imidazole (1.36 g imidazole) Adjust the buffer pH to 8.0 with NaOH solution.
Elution Buffer	1 L	50 mM NaH ₂ PO ₄ (7.80 g NaH ₂ PO ₄ ·2H ₂ O) 300 mM NaCl (17.54 g NaCl) 250 mM imidazole (17.0 g imidazole) Adjust the buffer pH to 8.0 with NaOH solution.

Table 4. Recommended buffer for his-tagged protein purification under denaturing conditions, pH elution

Name	Volume	Ingredient
Lysis Buffer	1 L	8 M Urea (480.50 g Urea) 100 mM NaH ₂ PO ₄ (15.60 g NaH ₂ PO ₄ ·2H ₂ O) 100 mM Tris·HCl (15.76 g Tris·HCl) Adjust the buffer pH to 8.0 with HCl solution
Wash Buffer	1 L	8 M Urea (480.50 g Urea) 100 mM NaH ₂ PO ₄ (15.60 g NaH ₂ PO ₄ ·2H ₂ O) 100 mM Tris·HCl (15.76 g Tris·HCl) Adjust the buffer pH to 6.3 with HCl solution
Elution Buffer	1 L	8 M Urea (480.50 g Urea) 100 mM NaH ₂ PO ₄ (15.60 g NaH ₂ PO ₄ ·2H ₂ O) 100 mM Tris·HCl (15.76 g Tris·HCl) Adjust the buffer pH to 4.5 with HCl solution





Table 5. Recommended buffer for his-tagged protein purification under denaturing conditions, imidazole elution

Name	Volume	Ingredient
Lysis Buffer	1 L	8 M Urea (480.50 g Urea) 50 mM NaH ₂ PO ₄ (7.80 g NaH ₂ PO ₄ ·2H ₂ O) 300 mM NaCl (17.54 g NaCl) 10 mM imidazole (0.68 g imidazole) Adjust the buffer pH to 8.0 with NaOH solution.
Wash Buffer	1 L	8 M Urea (480.50 g Urea) 50 mM NaH ₂ PO ₄ (7.80 g NaH ₂ PO ₄ ·2H ₂ O) 300 mM NaCl (17.54 g NaCl) 20 mM imidazole (1.36 g imidazole) Adjust the buffer pH to 8.0 with NaOH solution.
Elution Buffer	1 L	8 M Urea (480.50 g Urea) 50 mM NaH ₂ PO ₄ (7.80 g NaH ₂ PO ₄ ·2H ₂ O) 300 mM NaCl (17.54 g NaCl) 250 mM imidazole (17.0 g imidazole) Adjust the buffer pH to 8.0 with NaOH solution.

Note: 8 M Urea could also be replaced by 6 M Guanidine-HCl. Table 4 or Table 5 select one of them as the solution for the inclusion body purification.

2.2 Sample Preparation

2.2.1 Recombinant native protein expressed in *E.coli* or yeast

- 1) Single colonies were cultured in LB medium. According to the instruction, adding the inducers for a period of time.
- 2) Harvest cells from an appropriate volume of bacterial culture by centrifugation at 7,000rpm (7,500×g) for 10-15min at 4°C. Discard supernatant and determine weight of pellet. Resuspend pellet in 1:10 ration (w/v) in lysis buffer and add lysozyme (0.2-0.4 mg/ml cell paste, if the host cell containing pLysS or pLysE, it can be without lysozyme) and PMSF (1 mM/ml cell paste).
- 3) If the concentration of cell suspension is higher, it is recommended to add 10µg/ml RNase A and 5µg/ml DNase I. Sonicate the cell suspension/lysate on ice.
- 4) Centrifuge the homogenized lysate at 10,000 rpm for 20 to 30 min at 4°C to clarify sample. Save supernatant.

2.2.2 Native protein expressed in yeast, insect or mammalian cells

- 1) Harvest the cells from an appropriate volume of culture by centrifugation at 5,000 rpm for 10-15 min at 4°C. Save supernatant. If the supernatant without EDTA, histidine and reductant, it can be purified directly, otherwise it need dialysis to 1×PBS under 4°C.
- 2) For a large volume of supernatant, it need precipitation by adding ammonium sulfate and dialysis to 1×PBS under 4°C.

2.2.3 Inclusion bodies from *E.coli*

- 1) Harvest cells from an appropriate volume of bacterial culture by centrifugation at 7,000 rpm for 10-15min at 4°C. Discard supernatant and determine weight of pellet.
- 2) Resuspend pellet in 1:10 ration (w/v) with Lysis Buffer. Sonicate the cell suspension/lysate on ice.
- 3) Centrifuge the homogenized sample at 10,000 rpm or 20 min at 4°C to pellet the inclusion.
- 4) Resuspend pellet in 1:10 ration (w/v) with denaturing Lysis Buffer (containing 8 M urea). Sonicate, as needed, to redissolve the pellet.
- 5) Analyze the concentration of the target protein and continue with purification protocols under denaturing conditions.

2.3 Sample Purification

- 1) Fill the syringe or pump tubing with distilled water. Remove the stopper and connect the column to the syringe (with the provided connector), or pump tubing, "drop to drop" to avoid introducing air into the column. Remove the snap-off end at the column outlet.
- 2) Wash the column with 3-5 column volumes of distilled water.
- 3) Equilibrate the column with at least 5 column volumes Lysis Buffer.
- 4) Load the pre-treated sample by using a Loop fitted to the connector or by pumping it onto the column.
- 5) Wash the column with 10-15 column volumes of wash buffer or until the absorbance reaches the baseline or no material appears.
- 6) Elute with elution buffer using a stepwise or linear gradient. For one-step elution, 5 column volumes are usually enough. Other volumes may be required if the interaction is difficult to break. Linear gradient elution can be used to separate proteins of different binding strengths with a small gradient, such as 20 column volumes or more.

2.4 Analysis

Identify the fractions containing the His-tagged protein. Use UV absorbance, SDS-PAGE, or western blot.





3. Cleaning-in-Place

A column used to purify protein from cell extract usually contains some soluble substances and cell debris that are nonspecifically absorbed onto the matrix.

Cleaning-in-Place eliminates material not removed by regeneration and prevents progressive buildup of contaminants. If the column is to be reused, these contaminants should be cleaned from the column, as they were not completely removed during the sample clarification steps.

Remove the strong hydrophobic binding protein, lipoprotein and lipid

Wash the column with 5-10 column volumes 30% isopropanol contacting for 15-20 min. Or you can choose the 2 column volumes acidic or alkaline solution containing detergents, for example, 0.1 M acetic acid solution contains 0.1-0.5% non-ionic detergent, contacting for 1-2 hours .

Finally wash the column with 10 column volumes distilled water.

Remove the proteins combined with ion interacting

Wash the column with 1.5 M NaCl solution contacting for 10-15 min.

Finally wash the column with 10 column volumes distilled water.

4. Regenerating the medium

In general, **Ni NTA Beads 6FF** may be used a number of times before it becomes necessary to recharge them with metal ions. When the back pressure is too high or the capacity significantly lower, it need to strip the metal ions and recharge the **Ni NTA Beads 6FF** as the following procedure.

Wash the column with one of the following solutions.

- 1) Rinse with 5 column volumes of distilled water;
- 2) 100 mM EDTA (pH 8.0), 5 column volumes;
- 3) Rinse with 10 column volumes of distilled water;
- 4) 0.5 M NaOH, 5 column volumes, contacting for 10-15 min;
- 5) Rinse with 10 column volumes of distilled water;
- 6) 100 mM NiSO₄, 3-5 column volumes;
- 7) Rinse with 10 column volumes of distilled water;

After regeneration, the medium can be used immediately, otherwise, it need to be suspended in an equal volume of 1×PBS containing 20% ethanol at 4°C.

5. Troubleshooting

Problem	Probable Cause	Solution
Back pressure exceeds 3 bar	Column is clogged	Cleaning in place (part 3). Increase the centrifugation speed or filtering the sample.
	Sample is too viscous	Increase sonication or add DNase I (5 µg/ml with 1mM Mg ²⁺ . Incubate on ice for 15 min. Dilute sample with Binding buffer.
No protein is eluted	Protein may be an inclusion body, not in the supernatant.	The lysate can be analyzed by electrophoresis for the presence of the target protein in the supernatant, and the inclusion body protein need to be purified according to the purification method of the inclusion body protein.
	Expression of target protein in extract is very low	Check expression level of protein by estimating the amount in the extract, flow through, elute fraction and pellet upon centrifugation. Or apply large sample volume.
	Target protein is found in the flow through	Reduce imidazole concentration in sample lysis and wash buffer. Or increase the pH of buffer.
	Protein degradation or purification cause the histidine tag to be	Operate at 4°C. Add protease inhibitors. Make a new construct with his-tag attached to other terminus.





	removed.	
His-tagged protein is not pure	Wash is not enough	Increase the volume of Wash Buffer.
	Association between the his tagged protein and protein contaminant.	Optimize the wash condition by adjusting the pH and imidazole concentration. Add an additional chromatography step, that is ion exchange, hydrophobic interaction or size exclusion.
The color of medium becomes shallow.	The nickel ions was stripped.	Chelate nickel ions according to the part 4.
The color of medium becomes brown	The buffer contains reducing agent.	Refer to Table 2 to reduce the concentration of reducing agent
Protein precipitates during purification	Temperature is too high	Perform the purification at 4°C.
	Aggregate formation	Add solubilization agents to the samples and buffers, for example 0.1% Triton X-100 , Tween-20 and ≤20% glycerol to maintain protein solubility.

6. Related Products

Product	Cat. No.	Size
Ni NTA Beads	SA004005	5 ml
	SA004025	25 ml
	SA004100	100 ml
	SA004500	500 ml
	SA00401L	1 L
	SA00410L	10 L
HisPur Ni NTA Kit	SA004K03	3 times
Ni NTA Beads 6FF	SA005005	5 ml
	SA005025	25 ml
	SA005100	100 ml
	SA005500	500 ml
	SA00501L	1 L
	SA00510L	10 L
HisCap 6FF	SA005C11	1×1 ml
	SA005C51	5×1 ml
	SA005C15	1×5 ml
	SA005C55	5×5 ml
	SA005CS	3×1 ml+1×5 ml

